


New Directions in Searching for the Dark Universe

**Surjeet Rajendran,
UC Berkeley**

Dark Matter



A New Particle


Non gravitational interactions?

How do we detect them?

Weak effects. Need high precision


Precision Instruments

Impressive developments in the past two decades



$$\text{Magnetic Field} \lesssim 10^{-16} \frac{\text{T}}{\sqrt{\text{Hz}}}$$

(SQUIDs, atomic magnetometers)




$$\text{Accelerometers} \lesssim 10^{-13} \frac{\text{g}}{\sqrt{\text{Hz}}}$$

(atom and optical interferometers)

Rapid technological advancements

Use to detect new physics?

The Dark Matter Landscape



Fit in galaxy

Standard Model scale ~ 100 GeV

One Possibility: Same scale for Dark Matter?
Weakly Interacting Massive Particles (WIMPs)
Soon to hit solar neutrino floor

Other Generic Candidates: Axions, Massive Vector Bosons, Dark Blobs

How do we make progress?

Outline

1. Brief Theory Overview
2. Axion Detection with Nuclear Magnetic Resonance
3. Dark Photon Detection with Radios
4. Bosons with Accelerometers
5. Directional Detection of Dark Matter
6. Magnetic Bubble Chambers
7. Conclusions

Bosonic Dark Matter

Photons




$$\vec{E} = E_0 \cos(\omega t - \omega x)$$

Detect Photon by
measuring time varying
field

Dark Bosons

Early Universe:
Misalignment Mechanism



$$a(t) \sim a_0 \cos(m_a t)$$


Spatially uniform, oscillating field

$$m_a^2 a_0^2 \sim \rho_{DM}$$

Detect effects of oscillating dark matter field

Resonance possible. $Q \sim 10^6$ (set by $v \sim 10^{-3}$)

Today:
Random Field



Correlation length
 $\sim 1/(m_a v)$

Coherence Time
 $\sim 1/(m_a v^2)$
 $\sim 1 \text{ s (MHz}/m_a)$

What kind of Bosons?


Naturalness. Structure set by symmetries.

Spin 0


Spin 1

Axions or ultra weak coupling
Many UV theories


Anomaly free
Standard Model couplings




E&M




QCD




Spin




Higgs



Spin



E&M



Current

$$\left(\frac{a}{f_a} F \tilde{F} \right)$$

Current
Searches
($m_a \sim \text{GHz}$)

$$\left(\frac{a}{f_a} G \tilde{G} \right)$$

QCD
Axion

$$\left(\frac{\partial_\mu a}{f_a} \bar{N} \gamma^\mu \gamma_5 N \right)$$

General
Axions

$$(g \phi H^2)$$

Higgs Portal/
Relaxion

$$\left(\frac{F'_{\mu\nu}}{f_a} \bar{N} \sigma^{\mu\nu} N \right)$$

Dipole
moment

$$(\epsilon F' F) (g A'_\mu J_{B-L}^\mu)$$

Kinetic
Mixing

B-L

Dark Matter $\implies a = a_0 \cos(m_a t)$


This Talk

$$\text{Hz} \lesssim m_a \lesssim \text{GHz}$$

Observable Effects

What can the dark matter wind do?

What can a classical field do?



Cosmic Axion Spin Precession Experiment (CASPEr)

with

Dmitry Budker

Peter Graham

Micah Ledbetter

Alex Sushkov



PRX **4** (2014) arXiv: 1306.6089


PRD **88** (2013) arXiv: 1306.6088

PRD **84** (2011) arXiv: 1101.2691

CASPER: Axion Effects on Spin

General Axions

Neutron in
Neutron
Axion Wind



$$H_N \supset \frac{a}{f_a} \vec{v}_a \cdot \vec{S}_N$$

Spin rotates about
dark matter velocity


Effective time varying
magnetic field

$$B_{eff} \lesssim 10^{-16} \cos(m_a t) \text{ T}$$

QCD Axion

Neutron in
Neutron
QCD Axion Dark Matter

Measure Spin
Rotation,
detect Axion



QCD axion induces electric dipole moment
for neutron and proton

Dipole moment
along nuclear spin


$$\text{Oscillating dipole: } d \sim 3 \times 10^{-34} \cos(m_a t) \text{ e cm}$$

Apply electric field, spin rotates

Other light dark matter (e.g. dark photons) also
induce similar spin precession

CASPEr

Axion affects physics of nucleus, NMR is sensitive probe



Larmor frequency = axion mass \rightarrow resonant enhancement


SQUID measures resulting transverse magnetization

NMR well established technology, noise understood, similar setup to previous experiments

Example materials: LXe, ferroelectric PbTiO_3 , many others


CASPEr-General Axions

frequency (Hz)



~ year to scan one decade of frequency

CASPEr-QCD Axion



Verify signal with spatial coherence of axion field



Dark Photon Detection with a Radio

with

Peter Graham

Kent Irwin

Saptarshi Chaudhuri

Jeremy Mardon

Yue Zhao

arXiv: 1411.7382

Dark Photon Dark Matter

Many theories/vacua have additional, decoupled sectors, new U(1)'s

Natural coupling (dim. 4 operator): $\mathcal{L} \supset \varepsilon F F'$

mass basis:

$$\mathcal{L} = -\frac{1}{4} (F_{\mu\nu} F^{\mu\nu} + F'_{\mu\nu} F'^{\mu\nu}) + \frac{1}{2} m_{\gamma'}^2 A'_\mu A'^\mu - e J_{EM}^\mu (A_\mu + \varepsilon A'_\mu)$$


photon with small mass and suppressed couplings to all charged particles

**oscillating E' field
(dark matter)**

**can drive current
behind EM shield**


Dark Matter Radio Station

oscillating E' field
(dark matter)



Tunable resonant LC circuit
(a radio)

EXPECTED REACH



Parameters: volume $\sim 0.1 \text{ m}^3$, $T = 100 \text{ mK}$, $Q = 10^6$, 1 year

Dark Matter Detection with Accelerometers

with

Peter Graham

David Kaplan

Jeremy Mardon

William Terrano

B-L Dark Matter

Other than electromagnetism, only other anomaly free standard model current

$$\mathcal{L} = -\frac{1}{4} (F'_{\mu\nu} F'^{\mu\nu}) + \frac{1}{2} m_{\gamma'}^2 A'_\mu A'^\mu - g J_{B-L}^\mu A'_\mu$$

Protons, Neutrons, Electrons and Neutrinos are all charged

Electrically neutral atoms are charged under B-L

Force experiments constrain $g < 10^{-21}$

**oscillating E' field
(dark matter)**

can accelerate atoms

Force depends on net neutron number - violates equivalence principle. Dark matter exerts time dependent equivalence principle violating force!

The Relaxion

$$\mathcal{L} \supset (-M^2 + g\phi)|h|^2 + gM^2\phi + g^2\phi^2 + \dots + \Lambda^4 \cos \frac{\phi}{f}$$

Hierarchy problem solved through cosmic evolution - does not require any new physics at the LHC

ϕ is a light scalar coupled to higgs with small coupling g

$$\implies \frac{g\phi}{v} m_q \bar{q}q$$

$$\text{Dark matter } \phi \implies \phi = \phi_0 \cos(m_\phi(t - \vec{v} \cdot \vec{x}))$$

Time variation of masses of fundamental particles

$$\implies \text{force on atoms } \frac{g\nabla\phi}{v} m_q \sim \frac{gm_\phi\vec{v}}{v} m_q$$


Force violates equivalence principle. Time dependent equivalence principle violation!

Detection Options

Measure relative acceleration between different elements/isotopes.

Leverage existing EP violation searches and work done for gravitational wave detection

Torsion Balance




Force from dark matter causes torsion balance to rotate

Measure angle, optical lever arm enhancement

Atom Interferometer

Dark Matter




Differential free fall acceleration



Stanford Facility

Pulsar Timing Arrays



Pulsars are known to have stable rotation - can be used as clocks


Presently used to search for low frequency (100 nHz) gravitational waves.

Pulsar signal modulates due to gravitational wave passing between earth and the pulsar

Force by dark matter causes relative acceleration between Earth and Pulsar, leading to modulation of signal

Relaxion changes electron mass at location of Earth - changes clock comparison


Projected Sensitivities



Torsion Balance limited by fiber thermal noise

Atom interferometers by shot noise


Projected Sensitivities



Torsion Balance limited by fiber thermal noise

Atom interferometers by shot noise

The Dark Matter Landscape



Directional Detection of Dark Matter with Crystal Defects



(in progress)

with

Alex Sushkov and Nicholas Zobrist

Coherent Neutrino Scattering

Neutrinos and WIMPs have similar scattering topologies - rare, single particle collision with detector




Sun produces neutrinos.
Irreducible background.

Go beyond next generation?

Isotropic Dark Matter. Know location of Sun. Veto nuclear recoils coming from Sun's direction

Challenge: Big Target Mass. Need directional detection at solid state density.

Collision Aftermath



Tell-tale damage cluster well correlated with direction of initial ion,
localized within ~ 50 nm

Collision Aftermath


Tell-tale damage cluster well correlated with direction of initial ion,
localized within ~ 50 nm

Results of TRIM simulation, 30
keV initial ion


O(200 - 300) vacancies and
interstitials, lattice potential ~ 30
eV

Damage cascade well correlated
with direction of input ion

Need nano-scale measurement of
damage cascade



Nitrogen Vacancy Center in Diamond



Electronic levels sensitive to crystal environment ~ 50 nm scale


~ 1 per $(30 \text{ nm})^3$ of NV centers in bulk diamond demonstrated

Nano-scale measurements experimentally demonstrated. Active development of sensors by many groups around the world.

Can this be used for directional detection? What is the effect of the damage cascade on a NV center?

Note: similar phenomenology applies to F-centers of Metal Halides

Damage Cascade and NV Centers




● = Nitrogen ● = Carbon ● = Electron

Damage leads to strain in crystal.
Strain shifts transition line

$$\text{Strain: } \nabla u \propto \frac{1}{r^3} \times \mathcal{O}(100 - 300)$$

(Hooke's Law)



TRIM simulation of damage cascade - calculate strain using Hooke's law

NV center shift ~ 100 kHz @ 30 nm
Natural line width \sim kHz

Single NV center has sensitivity to cascade!

Detector Concept

Large detector, segments of thickness $\sim \text{mm}$

NV center density $\sim 1 \text{ per } (30 \text{ nm})^3$



Conventional WIMP scattering ideas (scintillation, ionization etc.) to localize interesting events

Expect few events/year that could be WIMP or neutrinos

Pull out segments of interest. Conventional schemes localize events to within mm

Micron-scale localization by simply shining light - damaged area will have measurable frequency shifts



For nano-scale resolution, apply external magnetic field gradient - hence need segmentation



Results

Take crystal. Grid of NV centers with density 1 per $(30 \text{ nm})^3$

Run ~ 1000 TRIM simulations, get cascade for each. Can grid distinguish direction (including head vs tail)?



More damage in tail vs head used for discrimination. Above 10 keV, efficiency $> 80\%$, false positive $< 4\%$

5 σ detection with few events!

Magnetic Bubble Chambers

(in progress)

with

Phil Bunting, Giorgio Gratta, Jeffrey Long and Tom Melia

The Dark Matter Landscape




What about this range?

Coherence time of signal too short for phase measurement to work. Energy deposition too small to be been using conventional WIMP calorimeters

Need amplification of deposited energy (meV - keV)

Challenge: Need large target mass. Rare dark matter event. Requires amplifier stability $>$ years

Concept




Consider magnet with all spins aligned

Spins now in metastable excited state with energy $\sim g \mu B$


Dark Matter collides, deposits heat. Causes meta-stable spin to flip

Spin flip releases stored Zeeman energy (exothermic). Released energy causes other spins to flip, leading to magnetic deflagration (burning) of material.




Amplifies deposited energy. Like a bubble chamber. Is this possible? Stability?

Single Molecular Magnets



Will not happen in a ferromagnet - spins are strongly coupled.

Need weak spin-spin coupling. But need large density - necessary for heat conduction. Can't use gas.




Organo-Metallic complexes.
Central metal complex surrounded by organic material.

Weak coupling between adjacent metal complexes - but still large density

Each molecule acts as an independent magnet

Recently discovered systems. Few 100 known examples. Can make large samples.
Magnetic deflagration experimentally observed and well studied in Manganese Acetate complexes

Magnetic Deflagration



System well described by 2 level Hamiltonian.
Two states separated by energy barrier.


Turn on magnetic field, metastable state decays to ground state through tunneling

$$\tau \propto \tau_0 \exp(U_{eff}/T)$$

Ultra-long lived state at low temperature - localized heating
rapidly decreases life-time, decay results in more energy release

Condition for Deflagration

Initially heat region of size λ to T



Deflagration occurs as long as we heat a sufficiently large region

U_{eff} and τ_0 sets the detector threshold. Short τ_0 and small U_{eff} means tiny energy deposit will sufficiently heat up material to trigger deflagration. Low threshold

Known examples with $\tau_0 \sim 10^{-13}$ s, $U_{\text{eff}} \sim 70$ K, enabling 0.01 eV thresholds



Detector Stability

High energy ($> \text{MeV}$) background from radio-active decays.

Detect MeV events using conventional means. Actual background at low energy very low - forward scattering of compton events

Problem: MeV events will constantly set off detector. Reset time vs operation time? Big problem for bubble chambers like COUPP

Expected background $\sim 1/(\text{m}^2 \text{ s})$. Initial detector size $\sim (10 \text{ cm})^3$ (kg mass), 1 background event $\sim 100 \text{ s}$



\vec{B}




With precision magnetometers, don't need entire crystal to flip

Within $\sim 10 \mu\text{s}$, flame $\sim 10 - 100 \mu\text{m}$. Visible with SQUID.

Shut off B, turn off fuel.
Deflagration stops. Lose $\sim (10 - 100 \mu\text{m})^3$ of volume every 100 s.

Potential Reach


$$\mathcal{L} = -\frac{1}{4} (F_{\mu\nu}F^{\mu\nu} + F'_{\mu\nu}F'^{\mu\nu}) + \frac{1}{2}m_{\gamma'}^2 A'_\mu A'^\mu - eJ_{EM}^\mu (A_\mu + \varepsilon A'_\mu)$$



Absorption obtained from photoabsorption. Exposure of 1 kg-year

Conclusions

The Dark Matter Landscape



Poor observational constraints on dark matter

Experiments under development can now search for dark matter particles with mass between 10^{-22} eV - 10^{-6} eV, using a variety of precision measurement tools

Explored concepts for WIMP directional detection in solid state densities and single molecular magnets for dark matter in the range 10^{-4} eV - GeV

Need to develop tools to cover full range of possibilities